



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

Draft genome sequences of two clostridium algidicarnis strains isolated from meat juice samples of chilled vacuum-packed lamb meat

Wambui, Joseph ; Cernela, Nicole ; Stevens, Marc J A ; Ghielmetti, Giovanni ; Stephan, Roger

Abstract: Clostridium algidicarnis causes blown-pack spoilage of vacuum-packed meat. Here, we report the draft genome sequences of C. algidicarnis strains CM002 and CM003, isolated from unspoiled chilled vacuum-packed lamb. The genome sequences of CM002 and CM003 comprise 2,950,326 and 2,870,995 bp, respectively, and each have a GC content of 30.1%

DOI: <https://doi.org/10.1128/mra.00983-20>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-200143>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Wambui, Joseph; Cernela, Nicole; Stevens, Marc J A; Ghielmetti, Giovanni; Stephan, Roger (2020). Draft genome sequences of two clostridium algidicarnis strains isolated from meat juice samples of chilled vacuum-packed lamb meat. Microbiology Resource Announcements, 9(45):e00983-20.

DOI: <https://doi.org/10.1128/mra.00983-20>



Draft Genome Sequences of Two *Clostridium algidicarnis* Strains Isolated from Meat Juice Samples of Chilled Vacuum-Packed Lamb Meat

 Joseph Wambui,^a Nicole Cernela,^a Marc J. A. Stevens,^a Giovanni Ghielmetti,^a  Roger Stephan^a

^aInstitute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland

ABSTRACT *Clostridium algidicarnis* causes blown-pack spoilage of vacuum-packed meat. Here, we report the draft genome sequences of *C. algidicarnis* strains CM002 and CM003, isolated from unspoiled chilled vacuum-packed lamb. The genome sequences of CM002 and CM003 comprise 2,950,326 and 2,870,995 bp, respectively, and each have a GC content of 30.1%.

Clostridium algidicarnis belongs to the group of psychrophilic and psychrotrophic clostridia causing blown-pack spoilage (BPS) in chilled vacuum-packed meat (1). Despite the role of this species in BPS, genomic studies of *C. algidicarnis* are limited. We report the draft genome sequences of the two nonhemolytic *C. algidicarnis* strains CM002 and CM003, isolated from meat juice samples from chilled vacuum-packed lamb (2).

The strains were isolated from two different meat juice samples following a multi-step sample preparation process involving ethanol (50% [vol/vol]; 1 h; 30°C) and lysozyme (4 mg/ml; 30 min; 37°C) treatment followed by anaerobic enrichment at 4°C in prereduced peptone-yeast-glucose-starch medium for 3 weeks. A loopful of the respective enriched sample was plated onto Columbia agar supplemented with 5% defibrinated sheep blood (CBA) and incubated further anaerobically at 4°C for 3 weeks. Single colonies were incubated anaerobically on CBA at 4°C for 2 weeks for purification. Anaerobic conditions were generated in anaerobic jars using Thermo Scientific Oxoid AnaeroGen sachets.

A DNA blood and tissue kit (Qiagen, Hombrechtikon, Switzerland) was used to extract genomic DNA from a single colony purified on the CBA. Illumina Nextera DNA Flex chemistry and a MiniSeq sequencer (Illumina, San Diego, CA, USA) were used for the preparation and sequencing of transposon-based libraries, respectively. The sequence reads were 1,335,527 and 755,507 bp for CM002 and CM003, respectively. For both genome sequences, paired-end sequence outputs of 150 to 300 bp were produced. Quality control was carried out using FastQC 0.11.7 (3) before assembly using the SPAdes 3.0-based software (4) Shovill 1.0.9 (<https://github.com/tseemann/shovill>). Contigs of >500 bp were selected for assembly. The assembled genomes of CM002 and CM003 comprise 2,950,326 and 2,870,995 bp, respectively. Each genome has a GC content of 30.1%. The number of contigs, genome coverage, contig N_{50} value, and contig L_{50} value for CM002 are 54, 45×, 185,888 bp, and 6, respectively, while for CM003, they are 56, 65×, 182,926 bp, and 7, respectively. CheckM (5) was used to assess the quality of both genomes. Default parameters were used for all software and Web servers.

Strain identification was carried out *in silico* using 16S-based identification tools (5, 6) and whole-genome sequence-based digital DNA-DNA hybridization (dDDH) (7) and average nucleotide identity (ANI) (8) against *C. algidicarnis* strains B3^T (GenBank accession no. [JNLN000000000](#)) and DSM 15099 ([PTIS000000000](#)). Comparative dDDH analyses for subspecies delimitation resulted in values ranging between 48.7 and 65.8%, which are

Citation Wambui J, Cernela N, Stevens MJA, Ghielmetti G, Stephan R. 2020. Draft genome sequences of two *Clostridium algidicarnis* strains isolated from meat juice samples of chilled vacuum-packed lamb meat. Microbiol Resour Announc 9:e00983-20. <https://doi.org/10.1128/MRA.00983-20>.

Editor Steven R. Gill, University of Rochester School of Medicine and Dentistry

Copyright © 2020 Wambui et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).

Address correspondence to Joseph Wambui, joseph.wambui@uzh.ch.

Received 21 August 2020

Accepted 19 October 2020

Published 5 November 2020

below the 79% threshold (7), indicating that strains CM002 and CM003 are phylogenetically distinct.

Gene annotation and prediction were carried out using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (9) and Rapid Annotations using Subsystems Technology (RAST) (10). Genes encoding proteolytic, lipolytic, and saccharolytic proteins were predicted, thus providing an insight into the metabolic features of these strains that might be involved in BPS.

Data availability. The whole-genome shotgun project for CM002 and CM003 has been deposited at DDBJ/ENA/GenBank under the accession no. [JACKWX000000000](#) and [JACKWW000000000](#), respectively. The versions described in this paper are [JACKWX010000000](#) and [JACKWW010000000](#), respectively. The raw sequencing reads have been deposited in the SRA under the accession no. [SRX8947912](#) and [SRX8947913](#), respectively.

REFERENCES

1. Silva AR, Paulo EN, Sant'Ana AS, Chaves RD, Massaguer PR. 2011. Involvement of *Clostridium gasigenes* and *C. aldicarnis* in "blown pack" spoilage of Brazilian vacuum-packed beef. *Int J Food Microbiol* 148:156–163. <https://doi.org/10.1016/j.jfoodmicro.2011.05.016>.
2. Wambui J, Püntener S, Corti S, Cernela N, Stephan R. 2020. Detection of psychrophilic *Clostridium* spp. causing "blown pack" spoilage in meat juice samples from chilled vacuum-packed beef and lamb meat imported from different countries to Switzerland. *J Food Prot* 83:56–59. <https://doi.org/10.4315/0362-028X.JFP-19-321>.
3. Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
4. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
5. Lee I, Chalita M, Ha S-M, Na S-I, Yoon S-H, Chun J. 2017. ContEst16S: an algorithm that identifies contaminated prokaryotic genomes using 16S RNA gene sequences. *Int J Syst Evol Microbiol* 67:2053–2057. <https://doi.org/10.1099/ijsem.0.001872>.
6. Yoon S-H, Ha S-M, Kwon S, Lim J, Kim Y, Seo H, Chun J. 2017. Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *Int J Syst Evol Microbiol* 67:1613–1617. <https://doi.org/10.1099/ijsem.0.001755>.
7. Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14:60. <https://doi.org/10.1186/1471-2105-14-60>.
8. Yoon S-H, Ha S-M, Lim J, Kwon S, Chun J. 2017. A large-scale evaluation of algorithms to calculate average nucleotide identity. *Antonie van Leeuwenhoek* 110:1281–1286. <https://doi.org/10.1007/s10482-017-0844-4>.
9. Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity GM, Kodira CD, Kyrpides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of standard operating procedures (SOPs) for (meta)genomic annotation. *OMICS* 12:137–141. <https://doi.org/10.1089/omi.2008.0017>.
10. Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, III, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5:8365. <https://doi.org/10.1038/srep08365>.